

ABELIAN LENGTH CATEGORIES OF STRONGLY UNBOUNDED TYPE

HENNING KRAUSE

ABSTRACT. We discuss the notion of strongly unbounded type for abelian length categories; this is closely related to the Second Brauer–Thrall Conjecture for artin algebras. A new ingredient is the space of characters in the sense of Crawley-Boevey.

1. INTRODUCTION

The Second Brauer–Thrall Conjecture says that over infinite fields every finite dimensional algebra of infinite representation type is of strongly unbounded type [10]. In this note we take a fresh look at the conjecture. We consider abelian length categories which are Hom-finite over some commutative ring and discuss the analogue of ‘strongly unbounded type’ in this slightly more general setting. We use Crawley-Boevey’s theory of characters [5, 6, 7] and further ideas from [9, 11].

A character for an abelian length category \mathcal{C} is an integer valued function $\text{Ob } \mathcal{C} \rightarrow \mathbb{N}$ which is compatible with the exact structure (Definition 3.1). It turns out that the collection $\text{Sp } \mathcal{C}$ of irreducible characters carries a rich structure. It is a topological space via Ziegler’s topology [20], and we investigate its basic properties. For instance, the Second Brauer–Thrall Conjecture amounts to the fact that $\text{Sp } \mathcal{C}$ is discrete if and only if \mathcal{C} has only finitely many isomorphism classes of indecomposable objects (Theorem 3.4). A crucial property is the compactness of $\text{Sp } \mathcal{C}$. There is an elegant proof of the Second Brauer–Thrall Conjecture when $\text{Sp } \mathcal{C}$ is compact. On the other hand, we point out an obstruction for the compactness of $\text{Sp } \mathcal{C}$ in terms of certain minimal closed subsets of $\text{Sp } \mathcal{C}$ (Proposition 5.1). This is illustrated by looking at modules over hereditary artin algebras (Corollary 5.3). The closed subsets of $\text{Sp } \mathcal{C}$ we look at correspond to subcategories of \mathcal{C} which are closed under subobjects; their relevance became apparent in recent work [14, 17]. Here, we show that each irreducible character is determined by such a closed subset (Corollary 4.4).

At this stage, not much seems to be known about the space $\text{Sp } \mathcal{C}$. So we end this note with a list of open problems and hope to stimulate further progress.

2. ARTIN ALGEBRAS

The following conjecture is the modification of the *Second Brauer–Thrall Conjecture* suggested by Crawley-Boevey in [4, §1.6]. Recall that the *endolength* of a module M is the length of M when viewed as a module over its endomorphism ring.

Conjecture 2.1 (Brauer–Thrall II). *Let A be an artin algebra of infinite representation type. Then for some $n \in \mathbb{N}$ there are infinitely many non-isomorphic indecomposable A -modules which are of endolength n and of finite length over A .*

Without going into details, let us mention that this conjecture has been established for artin algebras over a field k when k is algebraically closed [1, 3], and more generally when k is perfect [2].

3. ABELIAN LENGTH CATEGORIES

Let k be a commutative ring and \mathbf{C} a k -linear abelian length category such that each morphism set in \mathbf{C} has finite length as a k -module. Suppose also that \mathbf{C} has only finitely many non-isomorphic simple objects.

Definition 3.1 (Crawley-Boevey [7]). A *character* for \mathbf{C} is a function $\chi: \mathbf{Ob} \mathbf{C} \rightarrow \mathbb{N}$ satisfying the following:

- (1) $\chi(X \oplus Y) = \chi(X) + \chi(Y)$ for all $X, Y \in \mathbf{Ob} \mathbf{C}$, and
- (2) $\chi(X) + \chi(Z) \geq \chi(Y)$ for each exact sequence $X \rightarrow Y \rightarrow Z \rightarrow 0$ in \mathbf{C} .

A character $\chi \neq 0$ is *irreducible* if χ cannot be written as a sum of two non-zero characters. Note that any character can be written in a unique way as a finite sum of irreducible characters [6]. The *degree* of a character χ is

$$\deg \chi = \sum_S \chi(S)$$

where S runs through a representative set of simple objects in \mathbf{C} .

We denote by $\mathbf{Sp} \mathbf{C}$ the set of irreducible characters for \mathbf{C} . Fix a morphism $\alpha: X \rightarrow Y$ in \mathbf{C} and $n \in \mathbb{N}$. For each character χ set

$$\chi(\alpha) = \chi(X) - \chi(Y) + \chi(\text{Coker } \alpha)$$

and let

$$\mathbf{U}_\alpha = \{\chi \in \mathbf{Sp} \mathbf{C} \mid \chi(\alpha) \neq 0\} \quad \text{and} \quad \mathbf{V}_{\alpha,n} = \{\chi \in \mathbf{Sp} \mathbf{C} \mid \chi(\alpha) \leq n\}.$$

The following lemma describes a topology on $\mathbf{Sp} \mathbf{C}$; it is the analogue of Ziegler's topology on the isomorphism classes of indecomposable pure-injective modules over a ring [20].

Lemma 3.2. *The subsets of the form \mathbf{U}_α where α runs through all morphisms in \mathbf{C} form a basis of open subsets for a topology on $\mathbf{Sp} \mathbf{C}$. The subsets of the form $\mathbf{V}_{\alpha,n}$ are closed. Moreover,*

$$\mathbf{V}_n = \{\chi \in \mathbf{Sp} \mathbf{C} \mid \deg \chi \leq n\}$$

is closed and compact.

Proof. Consider the abelian category $\mathbf{A} = \mathbf{Fp}(\mathbf{C}, \mathbf{Ab})$ of additive functors $F: \mathbf{C} \rightarrow \mathbf{Ab}$ into the category \mathbf{Ab} of abelian groups that admit a *presentation*

$$\mathbf{Hom}(Y, -) \longrightarrow \mathbf{Hom}(X, -) \longrightarrow F \longrightarrow 0.$$

A function $\chi: \mathbf{Ob} \mathbf{A} \rightarrow \mathbb{N}$ is called *additive* provided that $\chi(F) = \chi(F') + \chi(F'')$ if $0 \rightarrow F' \rightarrow F \rightarrow F'' \rightarrow 0$ is an exact sequence. Restricting a function $\chi: \mathbf{Ob} \mathbf{A} \rightarrow \mathbb{N}$ to $\mathbf{Ob} \mathbf{C}$ by setting $\chi(X) = \chi(\mathbf{Hom}(X, -))$ gives a bijection between the additive functions $\mathbf{Ob} \mathbf{A} \rightarrow \mathbb{N}$ and the characters $\mathbf{Ob} \mathbf{C} \rightarrow \mathbb{N}$. The inverse map takes a character χ for \mathbf{C} to the function $\mathbf{Ob} \mathbf{A} \rightarrow \mathbb{N}$ which again we denote by χ and which is defined by $\chi(F) = \chi(\alpha)$ when F is presented by a morphism $\alpha: X \rightarrow Y$ in \mathbf{C} ; this does not depend on the choice of α by Schanuel's Lemma.

Now the assertions follow from properties of additive functions on \mathbf{A} established in Lemmas B.5, B.6, and C.1 in [13]. For the compactness of \mathbf{V}_n one uses that \mathbf{C} has only finitely many simple objects. \square

Remark 3.3. To each irreducible character $\chi \in \mathbf{Sp} \mathbf{C}$ corresponds a simple object S_χ in some abelian quotient category of \mathbf{A} . The endomorphism ring of S_χ is a division ring; it is an interesting invariant of χ (see [13, Remark B.3]) but not relevant here.

The *character* and the *degree* of an object M in \mathbf{C} are defined by

$$\chi_M(-) = \text{length}_{\text{End}(M)} \text{Hom}(-, M) \quad \text{and} \quad \deg M = \deg \chi_M.$$

Note that χ_M is irreducible when M is indecomposable, by [7, Theorem 3.6]. A character of the form χ_M with M in \mathbf{C} is called *finite*.

Theorem 3.4. *Let \mathbf{C} be a k -linear Hom-finite abelian category having only finitely many non-isomorphic simple objects. Consider the following statements.*

- (1) *The number of isomorphism classes of indecomposable objects in \mathbf{C} is finite.*
- (2) *For every $n \in \mathbb{N}$ the number of non-isomorphic indecomposable objects in \mathbf{C} which are of degree n is finite.*
- (3) *Every irreducible character for \mathbf{C} is finite.*
- (4) *The space of irreducible characters is discrete.*

Then (2)–(4) are equivalent. Assuming in addition that Conjecture 2.1 holds, they are also equivalent to (1).

We need some preparations for the proof. Let $\mathbf{A} = \text{Lex}(\mathbf{C}^{\text{op}}, \mathbf{Ab})$ be the category of left exact functors $\mathbf{C}^{\text{op}} \rightarrow \mathbf{Ab}$. This is a Grothendieck abelian category and \mathbf{C} identifies via the Yoneda embedding taking X in \mathbf{C} to $\text{Hom}(-, X)$ with the full subcategory of finite-length objects in \mathbf{A} ; see [8, Chap. II] for details.

As before and following [7], we assign to each object M in \mathbf{A} its character χ_M (a function $\text{Ob } \mathbf{C} \rightarrow \mathbb{N} \cup \{\infty\}$), and M is by definition *endofinite* if the values of χ_M are finite.

Theorem 3.5 (Crawley-Boevey). *The assignment $M \mapsto \chi_M$ induces a bijection between the isomorphism classes of indecomposable endofinite objects in \mathbf{A} and the irreducible characters for \mathbf{C} .*

Proof. This is Theorem 3.6 in [7]; see also Proposition B.2 in [13]. \square

The next lemma shows that the finite irreducible characters form a dense subset of $\mathbf{Sp} \mathbf{C}$.

Lemma 3.6. *Let χ_M be an irreducible character for \mathbf{C} where M is an object in \mathbf{A} . Write $M = \text{colim}_i M_i$ as a filtered colimit of objects in \mathbf{C} and let \mathbf{U} be the set of irreducible characters of the form χ_N with N a direct summand of some M_i . Then χ_M belongs to the closure of \mathbf{U} .*

Proof. Let $\alpha: X \rightarrow Y$ be a morphism in \mathbf{C} and $F: \mathbf{A} \rightarrow \mathbf{Ab}$ the corresponding functor with presentation

$$\text{Hom}(Y, -) \longrightarrow \text{Hom}(X, -) \longrightarrow F \longrightarrow 0.$$

Observe that $\chi_M \in \mathbf{U}_\alpha$ iff $F(M) \neq 0$. We have $\text{colim}_i F(M_i) \xrightarrow{\sim} F(M)$, and therefore $\chi_M \in \mathbf{U}_\alpha$ implies $\chi_N \in \mathbf{U}_\alpha$ for some $\chi_N \in \mathbf{U}$. Thus χ_M belongs to the closure of \mathbf{U} . \square

Corollary 3.7. *The finite irreducible characters form a dense subset of $\mathbf{Sp} \mathbf{C}$.* \square

The following lemma identifies the isolated points in $\mathbf{Sp} \mathbf{C}$. Recall that a morphism $\phi: M \rightarrow N$ in \mathbf{C} is *left almost split* if ϕ is not a split monomorphism and every morphism $M \rightarrow N'$ in \mathbf{C} which is not a split monomorphism factors through ϕ .

Lemma 3.8. *Let $\chi \in \mathbf{Sp} \mathbf{C}$. Then $\{\chi\}$ is open if and only if there is a left almost split morphism $M \rightarrow N$ in \mathbf{C} with $\chi = \chi_M$.*

Proof. Suppose first that $\{\chi\} = U_\alpha$ for some morphism $\alpha: X \rightarrow Y$ in \mathbf{C} . It follows from Corollary 3.7 that $\chi = \chi_M$ for some indecomposable object M in \mathbf{C} . Choose a morphism $\beta: X \rightarrow M$ which does not factor through α and form the following pushout.

$$\begin{array}{ccc} X & \xrightarrow{\alpha} & Y \\ \beta \downarrow & & \downarrow \\ M & \longrightarrow & N \end{array}$$

It is easily checked that the morphism $M \rightarrow N$ is left almost split. Conversely, if $\alpha: M \rightarrow N$ is left almost split in \mathbf{C} , then $U_\alpha = \{\chi_M\}$. \square

Proof of Theorem 3.4. There are two cases. Suppose first that there is a simple object S that does not embed into an injective object in \mathbf{C} . Let E be an injective envelope of S in $\mathbf{Lex}(\mathbf{C}^{\text{op}}, \mathbf{Ab})$ and write $E = \bigcup_i E_i$ as the directed union of non-zero objects in \mathbf{C} . Note that E has degree one. Each E_i has a simple socle and is therefore indecomposable of degree one as well. On the other hand, the lengths of the E_i are unbounded. The object E yields an irreducible character χ_E for \mathbf{C} which lies in the closure of the χ_{E_i} , by Lemma 3.6. Thus the space $\mathbf{Sp} \mathbf{C}$ is indiscrete. It follows that each of (1)–(4) does not hold.

Now suppose that \mathbf{C} has enough injective objects. Let E be an injective cogenerator and A its endomorphism ring. Then A is an artin algebra and the functor $\text{Hom}(-, E)$ identifies \mathbf{C} with the category of finite-length A -modules. Note that each indecomposable object in \mathbf{C} is the source of a left almost split morphism.

(3) \Leftrightarrow (4): The finite characters in $\mathbf{Sp} \mathbf{C}$ are precisely the isolated points, by Lemma 3.8. Thus $\mathbf{Sp} \mathbf{C}$ is discrete iff all irreducible characters are finite.

(3) \Rightarrow (2): The set V_n of irreducible characters of degree at most n form a closed subset of $\mathbf{Sp} \mathbf{C}$ which is compact, by Lemma 3.2. Every character in V_n is finite and therefore isolated, by Lemma 3.8. Thus V_n is a finite set.

(2) \Rightarrow (3): Suppose there is an irreducible character χ which is not finite. Using the correspondence between irreducible characters for \mathbf{C} and indecomposable endofinite A -modules [5, §5.3], it follows that χ corresponds to an endofinite A -module which is not of finite length. This implies that there are infinitely many non-isomorphic indecomposable finite-length A -modules of some fixed endlength; see [5, §9.6]. For each A -module M we have

$$\deg M \leq \chi_M(A/\text{rad } A) \leq \chi_M(A) = \text{endol } M.$$

Thus for some $n \in \mathbb{N}$ the number of non-isomorphic indecomposable objects in \mathbf{C} which are of degree n is infinite.

(1) \Rightarrow (2): Clear.

(2) \Rightarrow (1): This follows from Conjecture 2.1, using that the degree of an A -module is bounded by its endlength. \square

Remark 3.9. The original notion of ‘strongly unbounded type’ asks for infinitely many $n \in \mathbb{N}$ such that the number of non-isomorphic indecomposable objects of length n is infinite [10]. We cannot expect to have infinitely many such n when *length* is replaced by *degree* as in Theorem 3.4. To see this, consider the category of finitely generated torsion modules over any discrete valuation domain: all indecomposables are of degree one.

4. SUBOBJECT-CLOSED SUBCATEGORIES

In this section we fix a k -linear Hom-finite abelian category \mathcal{C} . A full additive subcategory $\mathcal{D} \subseteq \mathcal{C}$ is called *subobject-closed* if each subobject of an object in \mathcal{D} belongs to \mathcal{D} . Note that in this case the inclusion $\mathcal{D} \rightarrow \mathcal{C}$ has a left adjoint; it takes an object $X \in \mathcal{C}$ to X/U where $U \subseteq X$ is the minimal subobject with $X/U \in \mathcal{D}$. Thus \mathcal{D} is a category having cokernels and we can define characters $\text{Ob } \mathcal{D} \rightarrow \mathbb{N}$ as before.

Lemma 4.1. *Let $\mathcal{D} \subseteq \mathcal{C}$ be a subobject-closed subcategory and $p: \mathcal{C} \rightarrow \mathcal{D}$ be the left adjoint of the inclusion. The map $\text{Sp } \mathcal{D} \rightarrow \text{Sp } \mathcal{C}$ sending χ to $\chi \circ p$ identifies $\text{Sp } \mathcal{D}$ with the closure of $\{\chi_M \in \text{Sp } \mathcal{C} \mid M \in \mathcal{D}\}$.*

Proof. Adapt the proof of Proposition 2.2 in [14]. \square

A subset $\mathcal{U} \subseteq \text{Sp } \mathcal{C}$ is said to be *subobject-closed* if $\mathcal{U} = \text{Sp } \mathcal{D}$ for some subobject-closed additive subcategory $\mathcal{D} \subseteq \mathcal{C}$.

The subobject-closed subcategories of \mathcal{C} form a set which is partially ordered by inclusion. An intersection $\bigcap_{\alpha} \mathcal{C}_{\alpha}$ of subobject-closed additive subcategories $\mathcal{C}_{\alpha} \subseteq \mathcal{C}$ is again subobject-closed. Moreover, adapting the proof of [14, Corollary 2.3], we have

$$\text{Sp}(\bigcap_{\alpha} \mathcal{C}_{\alpha}) = \bigcap_{\alpha} \text{Sp } \mathcal{C}_{\alpha}.$$

Thus we can define for each $\chi \in \text{Sp } \mathcal{C}$ the subobject-closed subset

$$\text{sub } \chi = \bigcap_{\chi \in \mathcal{U}} \mathcal{U}$$

where \mathcal{U} runs through all subobject-closed subsets $\mathcal{U} \subseteq \text{Sp } \mathcal{C}$.

Theorem 4.2. *Let χ, ψ be irreducible characters for \mathcal{C} . Then*

$$\text{sub } \chi = \text{sub } \psi \iff \chi = \psi.$$

The proof is based on the following lemma.

Lemma 4.3. *Let M be an indecomposable endofinite object in $\mathbf{A} = \text{Lex}(\mathcal{C}^{\text{op}}, \mathbf{Ab})$ and $\phi: M \rightarrow \coprod_I M$ a monomorphism. Then at least one component of ϕ is invertible.*

Proof. Consider in \mathbf{A} the colimit of the chain of monomorphisms

$$M \xrightarrow{\phi} \coprod_I M \xrightarrow{\coprod_I \phi} \coprod_{I^2} M \xrightarrow{\coprod_I \coprod_I \phi} \coprod_{I^3} M \xrightarrow{\coprod_I \coprod_I \coprod_I \phi} \dots$$

and write this as morphism $\bar{\phi}: M \rightarrow \coprod_J M$. Denote by E the endomorphism ring of M ; it is a local ring with $\bigcap_{n \geq 0} \text{rad } E = 0$. This follows from [8, Prop. IV.14] since M can be identified with an indecomposable injective object of a locally finite abelian category; see [13, Proposition B.2]. If each component of ϕ belongs to $\text{rad } E$, then each component of $\bar{\phi}$ belongs to $\bigcap_{n \geq 0} \text{rad } E$. Thus there is one component of $\bar{\phi}$ which is invertible. \square

Proof of Theorem 4.2. As before, we work in the category $\mathbf{A} = \text{Lex}(\mathcal{C}^{\text{op}}, \mathbf{Ab})$. For $M \in \mathbf{A}$ let $\text{sub } M$ denote the subobject-closed subcategory of \mathcal{C} consisting of all finite-length subobjects of finite coproducts of copies of M . Note that M is a filtered colimit of objects in $\text{sub } M$.

Now fix indecomposable endofinite objects M and N in \mathbf{A} . Then

$$\text{sub } \chi_M = \text{Sp}(\text{sub } M)$$

and therefore

$$\mathbf{sub} \chi_M \subseteq \mathbf{sub} \chi_N \iff \mathbf{sub} M \subseteq \mathbf{sub} N.$$

Suppose that $\mathbf{sub} \chi_M \subseteq \mathbf{sub} \chi_N$. Write $M = \operatorname{colim}_i M_i$ as a filtered colimit of finite-length objects and choose for each i a monomorphism $\alpha_i: M_i \rightarrow N_i$ into a finite coproduct of copies of N . Note that the coproducts of copies of N form a subcategory of \mathbf{A} which is closed under filtered colimits and products, since N is endofinite; see [7, §3] or [12, Corollary 10.5]. Thus the α_i induce a monomorphism

$$M = \operatorname{colim}_i M_i \longrightarrow \operatorname{colim}_i \left(\prod_{j \rightarrow i} N_j \right) = \prod_I N$$

for some index set I .

Now suppose that $\mathbf{sub} \chi_M = \mathbf{sub} \chi_N$. Thus there is also a monomorphism $N \rightarrow \prod_J M$ for some index set J . Composing these morphisms yields a monomorphism

$$\phi: M \longrightarrow \prod_I \prod_J M.$$

Each component of ϕ is an endomorphism of M that factors through a coproduct of copies of N , and it follows from Lemma 4.3 that at least one component is invertible. Thus $M \cong N$ and therefore $\chi_M = \chi_N$. \square

Corollary 4.4. *The set $\mathbf{Sp} \mathbf{C}$ is partially ordered by*

$$\chi \subseteq \psi \iff \mathbf{sub} \chi \subseteq \mathbf{sub} \psi.$$

In particular, each $\chi \in \mathbf{Sp} \mathbf{C}$ is uniquely determined by the set of finite characters in $\mathbf{sub} \chi$.

Proof. The first part is clear from Theorem 4.2. For the second part, observe that $\mathbf{sub} \chi$ is the closure of the subset of finite characters in $\mathbf{sub} \chi$ by Lemma 4.3. \square

Let us rephrase Theorem 4.2 for module categories, using the correspondence between characters and endofinite modules [5, §5.3]. For a module M , let $\mathbf{sub} M$ denote the category consisting of all finite-length submodules of finite direct sums of copies of M .

Corollary 4.5. *Two indecomposable endofinite modules M and N over an artin algebra are isomorphic if and only if $\mathbf{sub} M = \mathbf{sub} N$.* \square

5. COMPACTNESS

We keep the setting of the previous section and discuss the compactness of $\mathbf{Sp} \mathbf{C}$. More specifically, the fact that each irreducible character is determined by a subobject-closed subset of $\mathbf{Sp} \mathbf{C}$ raises the question when a subset of $\mathbf{Sp} \mathbf{C}$ is of the form $\mathbf{sub} \chi$ for some character χ . We have the following criterion.

Proposition 5.1. *Suppose that each indecomposable object in \mathbf{C} is the source of a left almost split morphism. Let $\mathbf{U} \subseteq \mathbf{Sp} \mathbf{C}$ be an infinite subobject-closed subset which is minimal with respect to this property. Then the following conditions are equivalent.*

- (1) \mathbf{U} is compact.
- (2) \mathbf{U} contains an infinite character.
- (3) \mathbf{U} is of the form $\mathbf{sub} \chi$ for some character χ .

We need the following lemma.

Lemma 5.2. *Let $\mathbf{U} \subseteq \mathbf{Sp} \mathbf{C}$ and $\chi \in \mathbf{Sp} \mathbf{C}$. Choose indecomposable endofinite objects $(M_i)_{i \in I}$ and M in $\mathbf{A} = \mathbf{Lex}(\mathbf{C}^{\text{op}}, \mathbf{Ab})$ such that $\mathbf{U} = \{\chi_{M_i} \mid i \in I\}$ and $\chi = \chi_M$. Then the following conditions are equivalent.*

- (1) *The character χ belongs to the closure of \mathbf{U} .*
- (2) *The object M belongs to the smallest subcategory of \mathbf{A} closed under products, filtered colimits, pure subobjects, and containing all M_i .*

Proof. This follows from Corollary 4.6 and Theorem 6.2 in [12]. \square

Proof of Proposition 5.1. As before, we work in the category $\mathbf{A} = \mathbf{Lex}(\mathbf{C}^{\text{op}}, \mathbf{Ab})$. We denote by \mathbf{D} the subobject-closed subcategory of \mathbf{C} such that $\mathbf{U} = \mathbf{Sp} \mathbf{D}$.

(1) \Rightarrow (2): Each finite character in $\mathbf{Sp} \mathbf{C}$ is isolated, by Lemma 3.8. It follows that \mathbf{U} contains an infinite character when \mathbf{U} is infinite and compact.

(2) \Rightarrow (3): Let $\chi \in \mathbf{U}$ be infinite. Then $\mathbf{sub} \chi \subseteq \mathbf{U}$, and the minimality of \mathbf{U} implies equality.

(3) \Rightarrow (1): Let $\mathbf{U} = \mathbf{sub} \chi$ with $\chi = \chi_M$ for some indecomposable endofinite object M in \mathbf{A} . It suffices to show that each infinite closed subset $\mathbf{V} \subseteq \mathbf{U}$ contains χ . Thus we choose such an infinite subset \mathbf{V} . The minimality of \mathbf{U} implies that the smallest subobject-closed subset of $\mathbf{Sp} \mathbf{C}$ containing \mathbf{V} coincides with \mathbf{U} . Write $M = \text{colim}_i M_i$ as a filtered colimit of objects in \mathbf{D} . It follows that for each i there is a monomorphism $M_i \rightarrow N_i$ with N_i a finite coproduct of indecomposable objects $N \in \mathbf{C}$ such that $\chi_N \in \mathbf{V}$. There are also monomorphisms $N_i \rightarrow P_i$ such that each P_i is a finite coproduct of copies of M , since $N_i \in \mathbf{D} = \mathbf{sub} M$. These monomorphisms induce a pair of monomorphisms

$$M = \text{colim}_i M_i \longrightarrow \text{colim}_i \left(\prod_{i \rightarrow j} N_j \right) \longrightarrow \text{colim}_i \left(\prod_{i \rightarrow j} P_j \right) = \prod_I M$$

for some index set I , as in the proof of Theorem 4.2. It follows from Lemma 4.3 that at least one component of this composite is invertible. Thus χ belongs to \mathbf{V} , by Lemma 5.2. We conclude that \mathbf{U} is compact. \square

There is an interesting consequence for the space of irreducible characters on the category of finite-length modules over a hereditary artin algebra. I am grateful to Claus Michael Ringel for providing the key observation.

Corollary 5.3. *Let A be a connected hereditary artin algebra and $\mathbf{mod} A$ be the category of finite-length A -modules. Then the space $\mathbf{Sp}(\mathbf{mod} A)$ is compact if and only if A is of finite or tame representation type.*

Proof. The assertion is clear from Theorem 3.4 when A is of finite representation type, and it follows by inspection of the Ziegler spectrum when A is tame, see [15, 16]. Now suppose that A is of wild representation type and denote by \mathbf{C} the category of preprojective modules; it is minimal among the subobject-closed subcategories of $\mathbf{mod} A$ which are of infinite type, by [17, Example 1]. There is no indecomposable endofinite A -module of infinite length which is a union of modules from \mathbf{C} , by [18]. Thus the corresponding subset $\mathbf{Sp} \mathbf{C}$ is not compact, by Proposition 5.1. It follows that $\mathbf{Sp}(\mathbf{mod} A)$ is not compact. \square

6. SOME OPEN PROBLEMS

Not much seems to be known about the space $\mathbf{Sp} \mathbf{C}$ of characters for a length category \mathbf{C} . For instance, we need a better understanding of the interplay between finite and infinite characters. In this section we address some open problems. We fix a k -linear Hom-finite abelian category \mathbf{C} having only finitely many non-isomorphic simple objects.

The support of a character. It would be interesting to find for each irreducible character χ some appropriate set of finite irreducible characters *supporting* χ . More specifically, we ask the following.

Question 6.1. Is every irreducible character of degree n in the closure of a set of finite irreducible characters of degree at most n ?

Subobject-closed subsets. The lattice of subobject-closed subsets of $\mathbf{Sp}\, \mathbf{C}$ contains a lot of information. For instance, the space $\mathbf{Sp}\, \mathbf{C}$ itself embeds naturally into this lattice, by Theorem 4.2. This motivates the following question.

Question 6.2. When is a subobject-closed subsets of $\mathbf{Sp}\, \mathbf{C}$ of the form $\mathbf{sub}\, \chi$ for some irreducible character χ ?

There is a compactness result for subobject-closed subcategories which is due to Ringel [17]; his proof uses properties of the Gabriel–Roiter measure. An alternative proof in [14] is based on the compactness of the Ziegler spectrum.

A full additive subcategory $\mathbf{D} \subseteq \mathbf{C}$ is of *infinite type* if there are infinitely many isomorphism classes of indecomposable objects in \mathbf{D} .

Proposition 6.3. *Suppose that each indecomposable object in \mathbf{C} is the source of a left almost split morphism. Then each subobject-closed additive subcategory of \mathbf{C} that is of infinite type contains one which is minimal among all subobject-closed additive subcategories of infinite type.*

Proof. Adapt the proof of Corollary 4.3 in [14]. □

From this proposition it follows that each infinite subobject-closed subset of $\mathbf{Sp}\, \mathbf{C}$ contains a minimal one.

Question 6.4. When does a minimal infinite subobject-closed subset of $\mathbf{Sp}\, \mathbf{C}$ contain an infinite character?

Note that such a minimal infinite subobject-closed subset with an infinite character χ is necessarily of the form $\mathbf{sub}\, \chi$, by Proposition 5.1.

Remark 6.5. Let $\mathbf{D} \subseteq \mathbf{C}$ be a minimal subobject-closed subcategory of infinite type. Then for each $n \in \mathbb{N}$ the number of isomorphism classes of indecomposable objects of length n in \mathbf{D} is finite; see [17, Theorem 2].

Compactness. When $\mathbf{Sp}\, \mathbf{C}$ is compact, one gets an elegant proof of the Second Brauer–Thrall Conjecture, because an infinite space cannot be discrete; see Theorem 3.4. This observation motivates the following question.

Question 6.6. When is $\mathbf{Sp}\, \mathbf{C}$ compact?

Even when $\mathbf{Sp}\, \mathbf{C}$ is not compact, we can ask for suitable subsets which are compact. Here is one possible question.

Question 6.7. When is a subset of the form $\mathbf{sub}\, \chi$ compact?

Partial orders on the set of characters. The set of characters $\mathbf{Ob}\, \mathbf{C} \rightarrow \mathbb{N}$ is partially ordered by

$$\chi \leq \psi \iff \chi(X) \leq \psi(X) \text{ for all } X \in \mathbf{Ob}\, \mathbf{C}.$$

Observe that for each character ψ the set

$$\mathbf{U}_\psi = \{\chi \in \mathbf{Sp}\, \mathbf{C} \mid \chi \leq \psi\}$$

is closed and compact; this follows from Lemma 3.2.

From Theorem 4.2 it follows that the inclusion relation for $\mathbf{sub}\, \chi$ provides another partial order on the set of irreducible characters.

Question 6.8. How are the partial orders \leq and \subseteq on $\mathrm{Sp}\,C$ related?

We refer to [14, 19] for more details on these partial orders.

REFERENCES

- [1] R. Bautista, On algebras of strongly unbounded representation type. *Comment. Math. Helv.* **60** (1985), no. 3, 392–399.
- [2] R. Bautista and L. Salmerón, On discrete and inductive algebras, in *Representations of algebras and related topics*, 17–35, Fields Inst. Commun., 45 Amer. Math. Soc., Providence, RI, 2005.
- [3] K. Bongartz, Indecomposables are standard, *Comment. Math. Helv.* **60** (1985), no. 3, 400–410.
- [4] W. Crawley-Boevey, Tame algebras and generic modules, *Proc. London Math. Soc.* (3) **63** (1991), no. 2, 241–265.
- [5] W. Crawley-Boevey, Modules of finite length over their endomorphism rings, in *Representations of algebras and related topics (Kyoto, 1990)*, 127–184, London Math. Soc. Lecture Note Ser., 168 Cambridge Univ. Press, Cambridge, 1992.
- [6] W. Crawley-Boevey, Additive functions on locally finitely presented Grothendieck categories, *Comm. Algebra* **22** (1994), no. 5, 1629–1639.
- [7] W. Crawley-Boevey, Locally finitely presented additive categories, *Comm. Algebra* **22** (1994), no. 5, 1641–1674.
- [8] P. Gabriel, Des catégories abéliennes, *Bull. Soc. Math. France* **90** (1962), 323–448.
- [9] I. Herzog, The Ziegler spectrum of a locally coherent Grothendieck category, *Proc. London Math. Soc.* (3) **74** (1997) 503–558.
- [10] J. P. Jans, On the indecomposable representations of algebras, *Ann. of Math.* (2) **66** (1957), 418–429.
- [11] H. Krause, Generic modules over Artin algebras, *Proc. London Math. Soc.* (3) **76** (1998), no. 2, 276–306.
- [12] H. Krause, Exactly definable categories, *J. Algebra* **201** (1998), no. 2, 456–492.
- [13] H. Krause, Cohomological length functions, arXiv:1209.0540.
- [14] H. Krause and M. Prest, The Gabriel-Roiter filtration of the Ziegler spectrum, *Quart. J. Math.*, doi:10.1093/qmath/has020.
- [15] M. Prest, Ziegler spectra of tame hereditary algebras, *J. Algebra* **207** (1998), no. 1, 146–164.
- [16] C. M. Ringel, The Ziegler spectrum of a tame hereditary algebra, *Colloq. Math.* **76** (1998), no. 1, 105–115.
- [17] C. M. Ringel, Minimal infinite submodule-closed subcategories, *Bull. Sci. Math.* **136** (2012), no. 7, 820–830.
- [18] C. M. Ringel, Generic modules over wild hereditary artin algebras, in preparation.
- [19] K. Schmidt, The endofinite spectrum of a tame algebra, *J. Algebra* **279** (2004), no. 2, 771–790.
- [20] M. Ziegler, Model theory of modules, *Ann. Pure Appl. Logic* **26** (1984), no. 2, 149–213.

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT BIELEFELD, D-33501 BIELEFELD, GERMANY
E-mail address: hkrause@math.uni-bielefeld.de